



# Adequan® i.m. SINGLE DOSE

## Brand of Polysulfated Glycosaminoglycan (PSGAG)

Solution 500 mg/5 mL

For Intramuscular Use In Horses



**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**DESCRIPTION:** Each 5 milliliters of Adequan® i.m. contains 500 mg of Polysulfated Glycosaminoglycan and Water for Injection q.s. Sodium Hydroxide and/or Hydrochloric Acid added when necessary to adjust pH. Sodium Chloride may be added to adjust tonicity.

**PHARMACOLOGY:** Polysulfated Glycosaminoglycan is chemically similar to the glycosaminoglycans in articular cartilage matrix. PSGAG is a potent proteolytic enzyme inhibitor and diminishes or reverses the pathologic processes of traumatic or degenerative joint disease which result in a net loss of cartilage matrix components. PSGAG improves joint function by reducing synovial fluid protein levels and increasing synovial fluid hyaluronic acid concentration in traumatized equine carpal and hock joints.

**INDICATIONS:** Adequan® i.m. is recommended for the intramuscular treatment of non-infectious degenerative and/or traumatic joint dysfunction and associated lameness of the carpal and hock joints in horses.

**DOSAGE AND ADMINISTRATION:** The recommended dose of Adequan® i.m. in horses is 500 mg every 4 days for 28 days intramuscularly. The injection site must be thoroughly cleansed prior to injection. Do not mix Adequan® i.m. with other drugs or solvents.

**CONTRAINDICATIONS:** There are no known contraindications to the use of intramuscular Polysulfated Glycosaminoglycan.



# Adequan® i.m. SINGLE DOSE

**WARNINGS:** Do not use in horses intended for human consumption. Not for use in humans. Keep this and all medications out of the reach of children.

**PRECAUTIONS:** The safe use of Adequan® i.m. in horses used for breeding purposes, during pregnancy, or in lactating mares has not been evaluated.

**ANIMAL SAFETY:** Toxicity studies were conducted in horses. Doses as high as 2,500 mg were administered intramuscularly to 6 horses twice a week for 12 weeks. This dosage is 5 times the recommended dosage and 3 times the recommended therapeutic regimen. Clinical observations revealed no soreness or swelling at the injection site or in the affected joint. No animal had any clinical or laboratory evidence of toxicity.

**STORAGE CONDITIONS:** Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F) (See USP Controlled Room Temperature). Discard unused portion. Dispose of spent needles in accordance with all federal, state and local environmental laws.

**HOW SUPPLIED:** Adequan® i.m. solution, 500 mg/5 mL is available in 5 mL glass vials, packaged in boxes of 7.



**LUITPOLD PHARMACEUTICALS, INC.**

Animal Health Division

Shirley, NY 11967

(631) 924-4000

**(800) 458-0163**

Made in U.S.A.

IN 99501

Rev. 10/08

MG #7560

NADA #140-901, Approved by FDA



# Adequan<sup>®</sup> i.m. MULTI-DOSE

## Brand of Polysulfated Glycosaminoglycan (PSGAG)

Solution 100 mg/mL in 20 mL, 30 mL, and 50 mL Preserved Multi-Dose Vials

For Intramuscular Use In Horses Only. Not for Intra-Articular Use.



**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**DESCRIPTION:** Each mL contains Polysulfated Glycosaminoglycan 100 mg, Benzyl Alcohol 0.9% v/v as a preservative, and Water for Injection q.s. Sodium Hydroxide and/or Hydrochloric Acid added when necessary to adjust pH.

**PHARMACOLOGY:** Polysulfated Glycosaminoglycan is chemically similar to the glycosaminoglycans in articular cartilage matrix. PSGAG is a potent proteolytic enzyme inhibitor and diminishes or reverses the pathologic processes of traumatic or degenerative joint disease which result in a net loss of cartilage matrix components. PSGAG improves joint function by reducing synovial fluid protein levels and increasing synovial fluid hyaluronic acid concentration in traumatized equine carpal and hock joints.

**INDICATIONS:** Adequan<sup>®</sup> i.m. Multi-Dose is recommended for the intramuscular treatment of non-infectious degenerative and/or traumatic joint dysfunction and associated lameness of the carpal and hock joints in horses.

**DOSAGE AND ADMINISTRATION:** The recommended dose of Adequan<sup>®</sup> i.m. Multi-Dose in horses is 500 mg every 4 days for 28 days intramuscularly. The injection site must be thoroughly cleansed prior to injection. Do not mix Adequan<sup>®</sup> i.m. Multi-Dose with other drugs or solvents.

**CONTRAINDICATIONS:** There are no known contraindications to the use of intramuscular Polysulfated Glycosaminoglycan.

**WARNINGS:** Do not use in horses intended for human consumption. Not for use in humans. Keep this and all medications out of the reach of children.

# Adequan® i.m. MULTI-DOSE



**PRECAUTIONS:** The safe use of Adequan® i.m. Multi-Dose in horses used for breeding purposes, during pregnancy, or in lactating mares has not been evaluated.

**SAFETY AND EFFICACY:** Safety and efficacy studies utilizing Adequan® i.m. Multi-Dose were not performed. Adequan® i.m. Multi-Dose was approved based on the conclusion that the safety and effectiveness of Adequan® i.m. Multi-Dose will not differ from that demonstrated for the original formulation of Adequan® i.m.

**ANIMAL SAFETY:** Animal Safety studies utilizing Adequan® i.m. Multi-Dose were not performed. Safety studies were conducted in horses using the single dose formulation. Doses as high as 2,500 mg were administered intramuscularly to 6 horses twice a week for 12 weeks. This dosage is 5 times the recommended dosage and 3 times the recommended therapeutic regimen. Clinical observations revealed no soreness or swelling at the injection site or in the affected joint. No animal had any clinical or laboratory evidence of toxicity.

**STORAGE CONDITIONS:** Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F) (See USP Controlled Room Temperature). Dispose of spent needles in accordance with all federal, state and local environmental laws.

**HOW SUPPLIED:** Adequan® i.m. Multi-Dose solution, 100 mg/mL is available in 20 mL, 30 mL and 50 mL glass vials, packaged in boxes of 1.



**LUITPOLD**

**LUITPOLD PHARMACEUTICALS, INC.**

Animal Health Division

Shirley, NY 11967

(631) 924-4000

**800-458-0163**

Made in U.S.A.

IN92901

Rev. 10/08

MG# 28493

NADA #140-901, Approved by FDA